

	Description	$\delta x_-$	$\delta y_-$	$\delta x_+$	$\delta y_+$	
(a)	$K$ -matrix 1st solution	-2	0.9	2	1	
(b)	$K$ -matrix 2nd solution	0.3	0.3	0.0	-0.5	
(c)	Remove slowly varying part in $P$ -vector	-0.7	0.2	0.5	0.6	
(d)	Generalised LASS → relativistic Breit–Wigner	2	3	-1	3	
(e)	Gounaris-Sakurai → relativistic Breit–Wigner	0.7	0.0	-0.1	0.8	
(f)	$K^*(1680)$	$m + \delta m$	-0.0	0.6	0.1	0.5
(g)		$m - \delta m$	-0.2	-0.5	0.2	-0.9
(h)		$\Gamma + \delta\Gamma$	-0.2	0.2	0.0	-0.2
(i)		$\Gamma - \delta\Gamma$	0.2	-0.1	0.5	-0.2
(j)	$f_2(1270)$	$m + \delta m$	-0.1	0.0	0.3	-0.2
(k)		$m - \delta m$	-0.0	0.1	0.2	-0.2
(l)		$\Gamma + \delta\Gamma$	-0.0	0.0	0.2	-0.2
(m)		$\Gamma - \delta\Gamma$	-0.1	0.0	0.2	-0.2
(n)	$K_2^*(1430)$	$m + \delta m$	0.3	0.2	0.2	-0.2
(o)		$m - \delta m$	-0.4	-0.2	0.3	-0.1
(p)		$\Gamma + \delta\Gamma$	-0.2	0.2	0.1	-0.2
(q)		$\Gamma - \delta\Gamma$	0.1	-0.1	0.3	-0.2
(r)	$r_{\text{BW}} = 0.0 \text{ GeV}^{-1}$	-2	0.7	-1	-0.3	
(s)	$r_{\text{BW}} = 3.0 \text{ GeV}^{-1}$	4	-2	4	2	
(t)	Add $K^*(1410)$ and $\rho(1450)$	-0.2	-0.2	0.3	-0.3	
(u)	Helicity formalism	-6	6	-8	2	
	Total model related	8	7	10	5	