
$$K_1(1270)^- \quad m_0 = 1285.03 \pm 0.47 \pm 1.06 \text{ MeV}/c^2; \Gamma_0 = 90.79 \pm 1.12 \pm 2.54 \text{ MeV}/c^2$$

Partial Fractions [%]

$|g|$

$\arg(g)[^\circ]$

$\rho(770)^0 K^-$

$50.66 \pm 0.84 \pm 2.21$

$\bar{K}^*(892)^0 \pi^-$

$25.25 \pm 0.56 \pm 1.80$

$0.520 \pm 0.008 \pm 0.024$

$-133.2 \pm 0.9 \pm 2.2$

$[K^- \pi^+]^{L=0} \pi^-$

$5.97 \pm 0.29 \pm 0.37$

$0.390 \pm 0.010 \pm 0.018$

$95.2 \pm 1.4 \pm 3.7$

$[\bar{K}^*(892)^0 \pi^-]^{L=2}$

$2.73 \pm 0.14 \pm 0.24$

$0.946 \pm 0.028 \pm 0.147$

$8.8 \pm 1.7 \pm 2.4$

$\omega(782) [\pi^+ \pi^-] K^-$

$1.73 \pm 0.11 \pm 0.16$

$0.208 \pm 0.008 \pm 0.011$

$33.0 \pm 2.1 \pm 12.5$
